

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
Please do not report the images to the
Image Problem Mailbox.

THIS PAGE BLANK (USPTO)



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 03 JUL 2000

WIPO

PCT

EPO - DG 1

02.06.2000

(70)

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99106656.4

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN
THE HAGUE,
LA HAYE, LE

10/04/00

THIS PAGE BLANK (USPTO)



Eur päisches
Patentamt

European
Patent Office

Office eur péen
des brevets

Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation

Anmeldung Nr.:
Application no.:
Demande n°: 99106656.4

Anmeldetag:
Date of filing: 01/04/99
Date de dépôt:

Anmelder:
Applicant(s):
Demandeur(s):
Stymne, Sten, Dr.
26831 Svalöv
SWEDEN

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:

Recombinant DNA molecules encoding enzymes of the biosynthetic pathway for the production of triacylglycerol

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:

C12N15/54, C12N9/10, C12N15/81, C12N15/82, C12N1/16, C12N5/10

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:

The original title of the application reads as follows : Recombinant DNA molecules encoding a new class of enzymes in the biosynthetic pathway for the production of triacylglycerol

THIS PAGE BLANK (USPTO)

RECOMBINANT DNA MOLECULES ENCODING A NEW CLASS OF ENZYMES
IN THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION OF
TRIACYLGLYCEROL

5

The present invention relates to the isolation, identification and characterization of recombinant DNA molecules encoding enzymes catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.

10

The invention further relates to novel type of enzymes and their encoding genes for transformation. More specifically, the invention relates to use of a type of genes encoding a not previously described type of enzymes hereinafter designated phospholipid:diacylglycerol acyltransferases (PDAT). This type of genes expressed alone in transgenic organisms will enhance the total amount of oil (triacylglycerols) produced in the cells. The PDAT genes, in combination with a gene for the synthesis of an uncommon fatty acid will, when expressed in transgenic organisms, enhance the levels of the uncommon fatty acids in the triacylglycerols.

15

20 There is considerable interest world-wide in producing chemical feedstock, such as fatty acids, for industrial use from renewable plant resources rather than non-renewable petrochemicals. This concept has broad appeal to manufacturers and consumers on the basis of resource conservation and provides significant opportunity to develop new industrial crops for agriculture.

25

There is a diverse array of unusual fatty acids in oils from wild plant species and these have been well characterised (see e.g. Badami & Patil, 1981). Many of these acids have industrial potential and this has led to interest in domesticating relevant plant species to enable agricultural production of particular fatty acids.

30

Development in genetic engineering technologies combined with greater understanding of the biosynthesis of unusual fatty acids now makes it possible to transfer genes coding for key enzymes involved in the synthesis of a particular fatty acid from a wild species into domesticated oilseed crops. In this way individual fatty acids can be produced in high purity and quantities at moderate costs.

35

In all crops like rape, sunflower, oilpalm etc., the oil (i.e. triacylglycerols) is the most valuable product of the seeds or fruits and other compounds like starch, protein, and fibre is regarded as by-products with less value. Enhancing the quantity of oil per weight basis at the expense of other compounds in oil crops would therefore increase the value of crop. If genes, regulating the allocation of reduced carbon into the production of oil can be up-regulated, the cells will accumulate more oil on the expense of other products. Such genes might not only be used in already high oil producing cells such as oil crops but could also induce significant oil production in moderate or low oil containing crops such as e.g. soy, oat, maize, potato, sugarbeets, and turnips as well as in micro-organisms.

Summary of the invention

Many of the unusual fatty acids of interest, e.g. medium chain fatty acids, hydroxy fatty acids, epoxy fatty acids and acetylenic fatty acids, have physical properties that are distinctly different from the common plant fatty acids. The present inventors have found that, in plant species naturally accumulating these uncommon fatty acids in their seed oil (i.e. triacylglycerol), these acids are absent, or present in very low amounts in the membrane (phospho)lipids of the seed. The low concentration of these acids in the membrane lipids is most likely a prerequisite for proper membrane function and thereby for proper cell functions. One aspect of the invention is that seeds of transgenic crops can be made to accumulate high amounts of uncommon fatty acids if these fatty acids are efficiently removed from the membrane lipids and channelled into seed triacylglycerols.

The inventors have identified a novel class of enzymes in plants catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the production of triacylglycerol and, presumably, lysophospholipids and that these enzymes (phospholipid:diacylglycerol acyltransferases abbreviated as PDAT) are involved in the removal of hydroxylated, epoxygenated fatty acids, and probably also other uncommon fatty acids such as medium chain fatty acids, from phospholipids in plants. Further, the same enzyme reaction was shown to be present in microsomal preparations from baker's yeast (*Saccharomyces cerevisiae*). A so called 'knock out' yeast mutant, disrupted in the respective gene was obtained and microsomal membranes from the mutant was shown to totally lack PDAT activity. Thus, it was proved that the disrupted gene encodes for a PDAT enzyme. In addition, two further genes from *Arabidopsis thaliana* were found

with an amino acid sequence having 42 % identity over 96 amino acids and an amino acid sequence having 47 % identity over 73 amino acids with the yeast enzyme.

5 In a first embodiment, this invention is directed to nucleic acid sequences that encode a PDAT. This includes sequences that encode biologically active PDATs as well as sequences that are to be used as probes, vectors for transformation or cloning intermediates. The PDAT encoding sequence may encode a complete or partial sequence depending upon the intended use. All or a portion of the genomic sequence, cDNA sequence, precursor PDAT or mature PDAT is intended.

10

In a different aspect, this invention relates to a method for producing a PDAT in a host cell or progeny thereof, including genetically engineered oil seeds, yeast and moulds or any other oil accumulating organism, via the expression of a construct in the cell. Cells containing a PDAT as a result of the production of the PDAT encoding sequence are also contemplated within the scope of the invention.

15

In a different embodiment, this invention also relates to methods of using a DNA sequence encoding a PDAT for increasing the oil-content within a cell.

20 Another aspect of the invention relates to the accommodation of high amounts of uncommon fatty acids in the triacylglycerol produced within a cell, by introducing a DNA sequence producing a PDAT that specifically removes these fatty acids from the membrane lipids of the cell and channel them into triacylglycerol. Plant cells having such a modification are also contemplated herein.

25

A PDAT of this invention includes any sequence of amino acids, such as a protein, polypeptide or peptide fragment obtainable from a microorganism, animal or plant source that demonstrates the ability to catalyse the production of triacylglycerol from a phospholipid and diacylglycerol under enzyme reactive conditions. By „enzyme reactive conditions“ is meant that any necessary conditions are available in an environment (e.g., such factors as temperature, pH, lack of inhibiting substances) which will permit the enzyme to function.

30

Other PDATs are obtainable from the specific sequences provided herein. Furthermore, 35 it will be apparent that one can obtain natural and synthetic PDATs, including modified amino acid sequences and starting materials for synthetic-protein modelling from the

exemplified PDATs and from PDATs which are obtained through the use of such exemplified sequences. Modified amino acid sequences include sequences that have been mutated, truncated, increased and the like, whether such sequences were partially or wholly synthesised. Sequences that are actually purified from plant preparations or are identical or encode identical proteins thereto, regardless of the method used to obtain the protein or sequence, are equally considered naturally derived.

Further, the nucleic acid probes (DNA and RNA) of the present invention can be used to screen and recover „homologous“ or „related“ PDATs from a variety of plant and microbial sources.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1.

PDAT activity in microsomal fractions of *S. cerevisiae*. Aliquots of lyophilised microsomal membranes (10 nmol phosphatidylcholine) from a wild type yeast (strain YN979) (lane 1-3, 3), a yeast mutant (strain B10280), disrupted in the YNROO8w gene (lane 4-6, 9) or the yeast mutant complemented with a single copy plasmid containing the PDAT gene (lane 7) were assayed for PDAT activity. 2 nmol sn-1-oleoyl-sn-2-[¹⁴C]-ricinoleoylphosphatidylcholine (lane 1-7) or sn-1-oleoyl-sn-2-[¹⁴C]-oleoylphosphatidylcholine (lane 8-9) and 5 nmol of dioleoyl-diacylglycerol (lane 2,5, 7-9) or rac-oleoyl-vernoleoyl-diacylglycerol (lane 3, 6) were added in benzene solution. The benzene was evaporated under N₂ (g) and 0.1 ml of 50 mM potassium phosphate, pH 7.2, was added. The suspension was thoroughly mixed and after 90 min at 30 °C the lipids were extracted in chloroform and separated on thin layer chromatography on silica gel 60 plates in hexan/diethylether/acetic acid (35:70:1.5). The radioactive lipids were visualised and quantified on the plates by electronic autoradiography (Instant imager, Packard, US). Abbreviations used: triacylglycerol, TAG, FA, fatty acid (i.e. oleic acid); 1-OH-TAG, monoricinoleoyl-triacylglycerol; 1-OH-1-epTAG, monoricinoleoyl-monovernoleoyl-triacylglycerol and OH-FA, ricinoleic acid.

Brief Description of the SEQ ID:

SEQ ID NO. 1: The amino acid sequence of the yeast ORF YNROO8w from *Saccharomyces cerevisiae*

SEQ ID NO. 2: Amino acid sequence of the region of the *Arabidopsis thaliana* genomic sequence (AC004557).

SEQ ID NO. 3: Amino acid sequence of the region of the *Arabidopsis thaliana* genomic sequence (AB006704).

- 5 SEQ ID NO. 4: The corresponding genomic DNA sequence of the amino acid sequence (SEQ ID NO. 5) of the yeast ORF YNROO8w from *Saccharomyces cerevisiae*.

The present invention can be essentially characterized by the following aspects:

- 10 1. Use of a PDAT gene (genomic clone or cDNA) for transformation.
2. Use of a DNA molecule according to item 1 wherein said DNA is used for transformation of any organism in order to be expressed in this organism and result in an active recombinant PDAT enzyme in order to increase oil content of the organism.
- 15 3. Use of a DNA molecule of item 1 wherein said DNA is used for transformation of any organism in order to prevent the accumulation of undesirable fatty acids in the membrane lipids.
4. Use according to item 1, wherein said PDAT gene is used for transforming transgenic oil accumulating organisms engineered to produce any uncommon fatty acid which is harmful if present in high amounts in membrane lipids, such as
20 medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.
5. Use according to item 1, wherein said PDAT gene is used for transforming organisms, and wherein said organisms are crossed with other oil accumulating
25 organisms engineered to produce any uncommon fatty acid which is harmful if present in high amounts in membrane lipids, comprising medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.
6. Use according to item 1, wherein the enzyme encoded by said PDAT gene or cDNA is coding for a PDAT with distinct acyl specificity.
- 30 7. Use according to item 1 wherein said PDAT encoding gene or cDNA, is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 30% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 1.
8. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived
35 from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an

amino acid sequence 40% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 1.

9. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 60% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 1
10. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 80% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 1.
11. Use according to claim 1 wherein said PDAT encoding gene or cDNA is derived from from plants or contain nucleotide sequences coding for an amino acid sequence 40% or more identical to the amino acid sequence of PDAT from *Arabidopsis thaliana* as presented in SEQ. ID. NO. 2 or 3.
12. Transgenic oil accumulating organisms comprising, in their genome, a PDAT gene transferred by recombinant DNA technology or somatic hybridization.
13. Transgenic oil accumulating organisms according to item 12 comprising, in their genome, a PDAT gene having specificity for substrates with particular uncommon fatty acid and the gene for said uncommon fatty acid.
14. Transgenic organisms according to item 12 or 13 which are selected from the group consisting of fungi, plants and animals.
15. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants.
16. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a storage organ specific promotor.
17. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a seed promotor.
18. Oils from organisms according to item 12 - 17.
19. A method for altering acyl specificity of a PDAT by alteration of the nucleotide sequence of a naturally occurring encoding gene and as a consequence of this alternation creating a gene encoding for an enzyme with novel acyl specificity.
20. A protein encoded by a DNA molecule according to item 1 or a functional fragment thereof.
21. A protein of item 20 designated phospholipid:diacylglycerol acyltransferase.

22. A protein of item 21 which has a distinct acyl specificity.
23. A protein of item 13 having the amino acid sequence as set forth in Fig. 1, 2 or 3 or an amino acid sequence with at least 30 % homology to said amino acid sequence.
24. A protein of item 23 isolated from *Saccharomyces cerevisiae*.

5

EXAMPLE 1

10 Determination of substrate utilisation by the acyl-CoA independent acyltransferase in the synthesis of triacylglycerols in microsomal preparations of developing castor bean endosperm and bakers yeast.

15 Microsomal membranes prepared from developing endosperm of castor bean (*Ricinus communis*) catalyse the selective transfer of ricinoleoyl-(12-hydroxy-9-octadecenoyl) and vernoloyl(12-epoxy-9-octadecenoyl) groups from both diacylglycerols and phosphatidylcholine into triacylglycerols. The substrate utilisation was investigated in castor bean microsomes by using radioactive *sn*-1-oleoyl-*sn*-2-[¹⁴C]ricinoleoyl-diacylglycerol (20 *sn*-2-[¹⁴C]-ricinoleoyl-diacylglycerol) or *sn*-1-oleoyl-*sn*-2-(¹⁴C)-ricinoleoyl-phosphatidylcholine (*sn*-2-(¹⁴C)-ricinoleoyl-phosphatidylcholine) together with different non-radioactive diacylglycerol.

The preparation of microsomal fractions of developing castor bean endosperm and freeze drying of the microsomes were performed in known manner. Assays with addition of diacylglycerol and phosphatidylcholine substrates were performed. The results 25 showed that if radioactive *sn*-2-[¹⁴C]ricinoleoyl-diacylglycerol was used as the only added substrate, 2.8 % of the radio-labelled ricinoleoyl chains were found in triacylglycerol with one ricinoleoyl group, 12.4 % of the radioactivity was found in triacylglycerol-species with two ricinoleoyl groups and only trace amounts were associated with triacylglycerol consisting of three ricinoleoyl groups. If incubations with 30 *sn*-2-[¹⁴C] ricinoleoyl-diacylglycerol were performed in a 1:4 (mol:mol) mixture with non-radioactive diacylglycerol species containing one vernoloyl group, the distribution of radioactivity between different molecular species of triacylglycerol changed only marginally compared to incubations with just radioactive substrate. Only 1.3% of the added ¹⁴C-labelled ricinoleoyl groups were metabolised into triacylglycerol species with 35 one ricinoleoyl and one vernoloyl group. Similarly, only marginal changes in the radioactive triacylglycerol molecular species was seen in incubations where *sn*-2-[¹⁴C]-

ricinoleoyl-diacylglycerol was mixed with non-labelled divernoloyl-diacylglycerol. However, by adding unlabelled diricinoleoyl-phosphatidylcholine together with sn-2-[^{14}C]-ricinoleoyl-diacylglycerol the radioactivity metabolised into the different triacylglycerol species were substantially altered.

5

Only trace amounts of radioactivity were detected in triacylglycerol species with one ricinoleoyl chain whereas the radioactivity in triacylglycerol with two ricinoleoyl groups were doubled as compared to incubations with only sn-2-[^{14}C]-ricinoleoyl-diacylglycerol added.

10

EXAMPLE 2

Transformation and expression of YNROORw gene in yeast

- 15 The yeast mutant (strain B 10280) disrupted in the YNROO8w gene, was transformed with the single copy plasmid pFL39 having the PDAT-gene (YNROO8w) under the control of the endogenous promotor region (583 bp 5' untranslated) inserted into the cloning cassette. The transformed yeast was pre-cultivated at 28 °C for 20 h in defined YNB medium without tryptophane added. Cells were harvested and re-suspended in
- 20 minimal medium (Meesters et al., 1996), supplemented with 16 g/l glycerol to the original volume of the growth culture. The culture was further incubated for 24 h after which cells were harvested by centrifugation. Microsomal fraction of the yeast was prepared as described in Example 1 above and was incubated in the presence of sn-2-[^{14}C]-ricinoleoyl-phosphatidylcholine (Fig 1, lane 7). This experiment clearly shows that
- 25 the PDAT activity could be restored by the expression of the YNROO8w gene in the mutant yeast strain B10280 normally lacking the PDAT-activity.

- The effect of the over-expression of the PDAT gene on the lipid accumulation was studied by transforming the wild-type yeast (strain SCY62) with a plasmid pJN92
- 30 containing the PDAT gene (YNR008w) under the control of a GALI-promotor. The transformed yeast was then cultivated at 28 °C in defined YBN medium lacking uracil. The expression of the PDAT gene was induced by the addition of 2 % (v/v) galactose after 10 hours growth and was further incubated for 18 hours. The yeast cells were harvested and the lipid content of the yeast was analysed by thin layer chromatography
- 35 and gas liquid chromatography. The total lipid content in the yeast with the over-expressed PDAT was 1.3 fold higher than in the control yeast transformed with an empty

plasmid pJN92. The expression of the PDAT gene had no effect on the growth rate as determined by optical density measurements. The elevated lipid content in the yeast transformed with PDAT as compared to the control yeast can be totally accounted for by an 80 % increase seen in the triacylglycerol content. The levels of the polar lipids and sterol esters were not significantly effected by the over-expression of the PDAT gene. Hence, these results clearly demonstrate the use of the PDAT gene in increasing the oil content in transgenic organisms.

SEQUENCE LISTING

<110> Stymne Dr., Sten

5 <120> RECOMBINANT DNA MOLECULES ENCODING A NEW CLASS OF
ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION
OF TRIACYLGLYCEROL

10 <130> stymne

<140>
<141>

<160> 5

15 <170> PatentIn Ver. 2.0

<210> 1
<211> 661
20 <212> PRT
<213> *Saccharomyces cerevisiae*

<400> 1
25 Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 30
30 Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 60
35 Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 80
40 Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 110
45 Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115 120 125
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130 135 140
50 Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145 150 155 160
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
55 165 170 175
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
180 185 190
60 Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
195 200 205

Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
 210 215 220
 5 Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
 225 230 235 240
 Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
 245 250 255
 10 Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
 260 265 270
 Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
 275 280 285
 15 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
 290 295 300
 20 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
 305 310 315 320
 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
 325 330 335
 25 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350
 30 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380
 35 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415
 40 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430
 45 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460
 50 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480
 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu
 485 490 495
 55 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met
 500 505 510
 60 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr
 515 520 525

Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp
 530 535 540
 5 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro
 545 550 555 560
 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser
 565 570 575
 10 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly
 580 585 590
 Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
 595 600 605
 15 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
 610 615 620
 20 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
 625 630 635 640
 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
 645 650 655
 25 Met Pro Phe Pro Met
 660
 30 <210> 2
 <211> 387
 <212> PRT
 <213> Arabidopsis thaliana
 35 <400> 2
 Val Gly Ser Asn Val Tyr Pro Leu Ile Leu Val Pro Gly Asn Gly Gly
 1 5 10 15
 Asn Gln Leu Glu Val Arg Leu Asp Arg Glu Tyr Lys Pro Ser Ser Val
 20 25 30
 40 Trp Cys Ser Ser Trp Leu Tyr Pro Ile His Lys Lys Ser Gly Gly Trp
 35 40 45
 45 Phe Arg Leu Trp Phe Asp Ala Ala Val Leu Leu Ser Pro Phe Thr Arg
 50 55 60
 Cys Phe Ser Asp Arg Met Met Leu Tyr Tyr Asp Pro Asp Leu Asp Asp
 65 70 75 80
 50 Tyr Gln Asn Ala Pro Gly Val Gln Thr Arg Val Pro His Phe Gly Ser
 85 90 95
 Thr Lys Ser Leu Leu Tyr Leu Asp Pro Arg Leu Arg Asp Ala Thr Ser
 100 105 110
 55 Tyr Met Glu His Leu Val Lys Ala Leu Glu Lys Lys Cys Gly Tyr Val
 115 120 125
 60 Asn Asp Gln Thr Ile Leu Gly Ala Pro Tyr Asp Phe Arg Tyr Gly Leu
 130 135 140

13

Ala Ala Ser Gly His Pro Ser Arg Val Ala Ser Gln Phe Leu Gln Asp
 145 150 155 160
 5 Leu Lys Gln Leu Val Glu Lys Thr Ser Ser Glu Asn Glu Gly Lys Pro
 165 170 175
 Val Ile Leu Leu Ser His Ser Leu Gly Gly Leu Phe Val Leu His Phe
 180 185 190
 10 Leu Asn Arg Thr Thr Pro Ser Trp Arg Arg Lys Tyr Ile Lys His Phe
 195 200 205
 Val Ala Leu Ala Ala Pro Trp Gly Gly Thr Ile Ser Gln Met Lys Thr
 210 215 220
 15 Phe Ala Ser Gly Asn Thr Leu Gly Val Pro Leu Val Asn Pro Leu Leu
 225 230 235 240
 Val Arg Arg His Gln Arg Thr Ser Glu Ser Asn Gln Trp Leu Leu Pro
 245 250 255
 20 Ser Thr Lys Val Phe His Asp Arg Thr Lys Pro Leu Val Val Thr Pro
 260 265 270
 25 Gln Val Asn Tyr Thr Ala Tyr Glu Met Asp Arg Phe Phe Ala Asp Ile
 275 280 285
 Gly Phe Ser Gln Gly Val Val Pro Tyr Lys Thr Arg Val Leu Pro Leu
 290 295 300
 30 Thr Glu Glu Leu Met Thr Pro Gly Val Pro Val Thr Cys Ile Tyr Gly
 305 310 315 320
 Arg Gly Val Asp Thr Pro Glu Val Leu Met Tyr Gly Lys Gly Gly Phe
 325 330 335
 35 Asp Lys Gln Pro Glu Ile Lys Tyr Gly Asp Gly Asp Gly Thr Val Asn
 340 345 350
 40 Leu Ala Ser Leu Ala Ala Leu Lys Val Asp Ser Leu Asn Thr Val Glu
 355 360 365
 Ile Asp Gly Val Ser His Thr Ser Ile Leu Lys Asp Glu Ile Ala Leu
 370 375 380
 45 Lys Glu Ile
 385
 50 <210> 3
 <211> 389
 <212> PRT
 <213> Arabidopsis thaliana
 55 <400> 3
 Leu Lys Lys Glu Gly Leu Lys Ala Lys His Pro Val Val Phe Ile Pro
 1 5 10 15
 60 Gly Ile Val Thr Gly Gly Leu Glu Leu Trp Glu Gly Lys Gln Cys Ala
 20 25 30

Asp Gly Leu Phe Arg Lys Arg Leu Trp Gly Gly Thr Phe Leu Cys Trp
 35 40 45
 5 Val Glu His Met Ser Leu Asp Asn Glu Thr Gly Leu Asp Pro Ala Gly
 50 55 60
 Ile Arg Val Arg Ala Val Ser Gly Leu Val Ala Ala Asp Tyr Phe Ala
 65 70 75 80
 10 Pro Gly Tyr Phe Val Trp Ala Val Leu Ile Ala Asn Leu Ala His Ile
 85 90 95
 Gly Tyr Glu Glu Lys Asn Met Tyr Met Ala Ala Tyr Asp Trp Arg Leu
 100 105 110
 15 Ser Phe Gln Asn Thr Glu Arg Asp Gln Thr Leu Ser Arg Met Lys Ser
 115 120 125
 20 Asn Ile Glu Leu Met Val Ser Thr Asn Gly Gly Lys Lys Ala Val Ile
 130 135 140
 Val Pro His Ser Met Gly Val Leu Tyr Phe Leu His Phe Met Lys Trp
 145 150 155 160
 25 Val Glu Ala Pro Ala Pro Leu Gly Gly Gly Gly Gly Pro Asp Trp Cys
 165 170 175
 Ala Lys Tyr Ile Lys Ala Val Met Asn Ile Gly Gly Pro Phe Leu Gly
 180 185 190
 30 Val Pro Lys Ala Val Ala Gly Leu Phe Ser Ala Glu Ala Lys Asp Met
 195 200 205
 35 Arg Met Thr Arg Thr Trp Asp Ser Thr Met Ser Met Leu Pro Lys Gly
 210 215 220
 Gly Asp Thr Ile Trp Gly Gly Leu Asp Trp Ser Pro Glu Leu Pro Asn
 225 230 235 240
 40 Ala Pro Glu Met Glu Ile Tyr Ser Leu Tyr Gly Val Gly Ile Pro Thr
 245 250 255
 Glu Arg Ala Tyr Val Tyr Lys Leu Asn Gln Ser Pro Asp Ser Cys Ile
 260 265 270
 45 Pro Phe Gln Ile Phe Thr Ser Ala His Glu Glu Asp Glu Asp Ser Cys
 275 280 285
 50 Leu Lys Ala Gly Val Tyr Asn Val Asp Gly Asp Glu Thr Val Pro Val
 290 295 300
 Leu Ser Ala Gly Tyr Met Cys Ala Lys Ala Trp Arg Gly Lys Thr Arg
 305 310 315 320
 55 Phe Asn Pro Ser Gly Ile Lys Thr Tyr Ile Arg Glu Tyr Asn His Ser
 325 330 335
 Pro Pro Ala Asn Leu Leu Glu Gly Arg Gly Thr Gln Ser Gly Ala His
 340 345 350
 60 Val Asp Ile Met Gly Asn Phe Ala Leu Ile Glu Asp Ile Met Arg Val

15

	355	360	365	
	Ala Ala Gly Gly Asn Gly Ser Asp Ile Gly His Asp Gln Val His Ser			
	370	375	380	
5	Gly Ile Phe Glu Trp			
	385			
10	<210> 4			
	<211> 1986			
	<212> DNA			
	<213> Saccharomyces cerevisiae			
15	<220>			
	<221> CDS			
	<222> (1)..(1983)			
20	<400> 4			
	atg ggc aca ctg ttt cga aga aat gtc cag aac caa aag agt gat tct		48	
	Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser			
	1 5 10 15			
25	gat gaa aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga		96	
	Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg			
	20 25 30			
30	aac cac att cat cat caa cag gga tta ggc cat aag aga aga agg ggt		144	
	Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly			
	35 40 45			
35	att agt ggc agt gca aaa aga aat gag cgt ggc aaa gat ttc gac agg		192	
	Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg			
	50 55 60			
40	aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg		240	
	Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu			
	65 70 75 80			
45	att ttc att ctt ggt gca ttc tta ggt gta ctt ttg ccg ttt agc ttt		288	
	Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe			
	85 90 95			
50	ggc gct tat cat gtt cat aat agc gat agc gac ttg ttt gac aac ttt		336	
	Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe			
	100 105 110			
55	gta aat ttt gat tca ctt aaa gtg tat ttg gat gat tgg aaa gat gtt		384	
	Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val			
	115 120 125			
60	ctc cca caa ggt ata agt tcg ttt att gat gat att cag gct ggt aac		432	
	Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn			
	130 135 140			
65	tac tcc aca tct tct tta gat gat ctc agt gaa aat ttt gcc gtt ggt		480	
	Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly			
	145 150 155 160			
70	aaa caa ctc tta cgt gat tat aat atc gag gcc aaa cat cct gtt gta		528	
	Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val			

16

	165	170	175	
5	atg gtt cct ggt gtc att tct acg gga att gaa agc tgg gga gtt att Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile 180 185 190	576		
10	gga gac gat gag tgc gat agt tct gcg cat ttt cgt aaa cgg ctg tgg Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp 195 200 205	624		
15	gga agt ttt tac atg ctg aga aca atg gtt atg gat aaa gtt tgt tgg Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp 210 215 220	672		
20	ttg aaa cat gta atg tta gat cct gaa aca ggt ctg gac cca ccg aac Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn 225 230 235 240	720		
25	ttt acg cta cgt gca gca cag gcc ttc gaa tca act gat tat ttc atc Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile 245 250 255	768		
30	gca ggg tat tgg att tgg aac aaa gtt ttc caa aat ctg gga gta att Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile 260 265 270	816		
35	ggc tat gaa ccc aat aaa atg acg agt gct gcg tat gat tgg agg ctt Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu 275 280 285	864		
40	gca tat tta gat cta gaa aga cgc gat agg tac ttt acg aag cta aag Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys 290 295 300	912		
45	gaa caa atc gaa ctg ttt cat caa ttg agt ggt gaa aaa gtt tgt tta Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu 305 310 315 320	960		
50	att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa tgg Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp 325 330 335	1008		
55	gtc gag gct gaa gcc cct ctt tac ggt aat ggt ggt cgt gcc tgg gtt Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val 340 345 350	1056		
60	aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg gcc Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly 355 360 365	1104		
65	gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr 370 375 380	1152		
70	att caa tta aat acg tta gcc atg tat ggt ttg gaa aag ttc ttc tca Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser 385 390 395 400	1200		
75	aga att gag aga gta aaa atg tta caa acg tgg ggt ggt ata cca tca Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser 405 410 415	1248		

	atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct	1296
	Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser	
	420 425 430	
5	tca gag gat gca ttg aat aac aac act gac aca tac ggc aat ttc att	1344
	Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile	
	435 440 445	
10	cga ttc gaa agg aat acg agc gat gct ttc aac aaa aat ttg aca atg	1392
	Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met	
	450 455 460	
15	aaa gac gcc att aac atg aca tta tcg ata tca cct gaa tgg ctc caa	1440
	Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln	
	465 470 475 480	
20	aga aga gta cat gag cag tac tcg ttc ggc tat tcc aag aat gaa gaa	1488
	Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu	
	485 490 495	
	gag tta aga aaa aat gag cta cac cac aag cac tgg tcg aat cca atg	1536
	Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met	
	500 505 510	
25	gaa gta cca ctt cca gaa gct ccc cac atg aaa atc tat tgt ata tac	1584
	Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr	
	515 520 525	
30	ggg gtg aac aac cca act gaa agg gca tat gta tat aag gaa gag gat	1632
	Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp	
	530 535 540	
35	gac tcc tct gct ctg aat ttg acc atc gac tac gaa agc aag caa cct	1680
	Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro	
	545 550 555 560	
40	gta ttc ctc acc gag ggg gac gga acc gtt ccg ctc gtg gcg cat tca	1728
	Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser	
	565 570 575	
	atg tgt cac aaa tgg gcc cag ggt gct tca ccg tac aac cct gcc gga	1776
	Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly	
	580 585 590	
45	att aac gtt act att gtg gaa atg aaa cac cag cca gat cga ttt gat	1824
	Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp	
	595 600 605	
50	ata cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctc ggc agc	1672
	Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser	
	610 615 620	
55	gcg gag ttg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat	1920
	Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp	
	625 630 635 640	
60	ctc gtc gag cca cgc caa ttg tct aat ttg agc cag tgg gtt tct cag	1968
	Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln	
	645 650 655	

atg ccc ttc cca atg taa
Met Pro Phe Pro Met
660

1986

5

<210> 5
<211> 661
<212> PRT
<213> Saccharomyces cerevisiae

10

<400> 5
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15

15

Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 30

Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45

20

Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 60

25

Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 80

Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95

30

Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 110

Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115 120 125

35

Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130 135 140

40

Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145 150 155 160

Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
165 170 175

45

Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
180 185 190

Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
195 200 205

50

Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
210 215 220

55

Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
225 230 235 240

Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
245 250 255

60

Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
260 265 270

Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
 275 280 285
 5 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
 290 295 300
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
 305 310 315 320
 10 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
 325 330 335
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350
 15 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365
 20 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400
 25 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430
 30 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445
 35 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480
 40 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu
 485 490 495
 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met
 500 505 510
 45 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr
 515 520 525
 Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp
 530 535 540
 50 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro
 545 550 555 560
 55 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser
 565 570 575
 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly
 580 585 590

20

Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
595 600 605

5 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
610 615 620

Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
625 630 635 640

10 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
645 650 655

Met Pro Phe Pro Met
15 660

Claims

1. A recombinant DNA molecule encoding an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol, wherein said enzyme comprises a amino sequence set forth in SEQ ID NO. 1, SEQ. ID. NO. 2 or SEQ ID NO. 3 or fragments thereof.
2. The DNA molecule of claim 1 wherein said enzyme is designated as phospholipid:diacylglycerol acyltransferase.
3. A vector comprising a DNA molecule of claim 1 or 2.
4. A vector of claim 3 further comprising a selectable marker gene.
5. A host cell containing a DNA molecule of claim 1 or 2.
6. The host cell of claim 5 which is a plant cell or yeast cell.
7. A process for the production of transgenic yeast cells, plant cells or plants comprising a) transforming a DNA molecule of claims 1 or 2 into plant cells or plants; and b) selecting of transformed plant cells or plants having an altered biosynthetic pathway in the production of triacylglycerol.
8. A process of claim 7 wherein the altered biosynthetic pathway is characterised by an increased or altered oil content.
9. A method of claim 7 wherein the altered biosynthetic pathway is characterised by the prevention of accumulation of undesirable fatty acids in the membrane lipids.
10. A protein encoded by a DNA molecule according to claim 1 or 2 or a functional fragment thereof

Fig 1

Abstract of the Disclosure

5 The present invention relates to the isolation, identification and characterization of recombinant DNA molecules encoding an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.

THIS PAGE BLANK (USPTO)